

References

- Perioperative Standards and Recommended Practices. AORN, Inc. 2008
- Centers for Disease Control and Prevention (CDC). Guideline for the Prevention of Intravascular Catheter-Related Infections. MMWR. 2002;51(RR-10)
- Centers for Disease Control and Prevention (CDC). Guideline for Prevention of Surgical Site Infection. 1999;24(No.4)
- Association of periOperative Registered Nurses (AORN). Inc. 2008. Guidance Statement: Fire Prevention in the Operating Room.
- Operating Room Nurses Association of Canada Recommended Standards (ORNAC). Guidelines and Position Statements for Perioperative Registered Nursing Practice, June 2009, 9th Edition.
- Centers for Medicare and Medicaid Services (CMS), Memorandum Summary; Use of Alcohol-based Skin Preparations in Anesthetizing Locations, 2007.
- Centers for Disease Control and Prevention (CDC), Guidelines for the Prevention of Intravascular Catheter-Related Infections, 2009 Draft.
- INS – Infusion Nurses Society, Infusion Nursing Standards of Practice, Lippincott Williams &Wilkins, 2006.
- Safer Healthcare Now, Getting Started Kit: Prevent Central Line Infections, Central Line Associated – Blood Stream Infections (CLA-BSI), How- To Guide, February 2009)
- Mayal G.M. “Hospital Epidemiology and Infection Control” Ed. William & Wilkin, 1996.
- Gruendemann, Barbara J., Stonehocker, Sandra, Infection Prevention in Surgical Setting, Saunders, 2001.
- 11th Ed. Berry & Kohn’s Operating Room Technique, Nancy Marie Phillips, Mosby, 2007.
- Foster L. Wallis M, Paterson B, et al. A descriptive study of peripheral intravenous catheters in patients admitted to a pediatric unit in one Australian hospital. J Inf Nurs 2002; 25(3).
- Aggett PJ, Cooper LV, Ellis SH, et al. Percutaneous absorption of chlorhexidine in neonatal cord care. Arch of Disease in Childhood 1981; 56:878-880.
- Bhutta A, Gilliam C, Honeycutt M, et al. Quality improvement report - reduction of blood stream infections associated with catheters in pediatric intensive care unit: stepwise approach. BMJ 2007; 334.
- Blumer JL, Husak MP, Wiltshire J, et al. Antibacterial efficacy and safety of neonatal bathing with Hibiclens. Pediatr Res 1982; 16(4):148A.
- Cowen J, Ellis SH, McAinsh J. Absorption of chlorhexidine from the intact skin of newborn infants. Arch Dis Childhood 1979; 54.
- Garland JS. A randomized trial comparing povidone-iodine to a chlorhexidine gluconate-impregnated dressing for prevention of central venous catheter infections in neonates. Pediatrics 2001; 107.
- Garland JS. Comparison of 10% povidone-iodine and 0.5% chlorhexidine gluconate for the prevention of peripheral intravenous catheter colonization in neonates: a prospective trial. Pediatr Infect Dis J 1995; 14.
- Jarrell N. Minimizing the Risk of Catheter Related Blood Stream Infections from Multiple Line Accesses in a Pediatric Cardiac Intensive Care Unit [poster]. Presented at the Annual Conference of the Association for Professionals in Infection Control and Epidemiology (APIC), June 24-28, 2007, San Jose, CA.

References CON'D

- Johnson J, Seeberg S, Kjellmer I. Blood concentration of chlorhexidine in neonates undergoing routine care with 4% chlorhexidine gluconate solution. *Acta Paediatr Scand* 1987;76.
- Kinirons B. Chlorhexidine versus povidone-iodine in preventing colonization of continuous epidural catheters in children. *Anesthesiology*. 2001;94(2).
- O'Neill J, Hosmer M, Challop R, et al. Percutaneous absorption potential of chlorhexidine in neonates. *Current Therap Res* 1982; 31(3).
- Aerls A, Nevolsen D, Renard S. *Soins de plaies*. Chapter 3 – Antiseptiques. 1998.
- Badrikian L, Boiko-Alaux V. *L'antiseptoguide: Guide d'utilisation des antiseptiques*, p. 21/52. Hygiene Hospitaliere, CHU de Clermont-Ferrand, France.
- Chaiyakunapruk, N. et al. Chlorhexidine Compared with Povidone-Iodine Solution for Vascular Catheter–Site Care: A Meta-Analysis. *American College of Physicians–American Society of Internal Medicine, Antiseptic for Catheter-Site Care: A Meta-Analysis*. June 2002; 136(11).
- Maki, Dennis G. et al. Prospective Randomized Trial of Povidone-Iodine, Alcohol, and Chlorhexidine for Prevention of Infection Associated with Central Venous and Arterial Catheters. *The Lancet*, 1991; 338.
- Small, Heather et al. Efficacy of Adding 2% (w/v) Chlorhexidine Gluconate to 70% (v/v) Isopropyl Alcohol for Skin Disinfection Prior to Peripheral Venous Cannulation. *Infection Control and Hospital Epidemiology*, October, 2008; 29(10).
- Pearce, B. et al. Efficacy of Clean v Sterile Surgical Prep Kits. *Association of periOperative Registered Nurses Journal (AORN)*. September, 1997; 66(3).
- Darouiche, Rabih O. et al. Chlorhexidine-Alcohol versus Povidone-Iodine for Surgical-Site Antisepsis. *The New England Journal of Medicine*, January, 2010; 362:18-26.
- Swensen, Brian R. et al. Effects of Preoperative Skin Preparation on Postoperative Wound Infection Rates: A Prospective Study of 3 Skin Preparation Protocols. *Infection Control and Hospital Epidemiology*, October, 2009; 30(10).
- Denton, Graham W. et al. *Disinfection, Sterilization and Preservation*. Chapter 15, Chlorhexidine; p. 321-336.
- Hibbard, John S. Analyses Comparing the Antimicrobial Activity and Safety of Current Antiseptic Agents: A Review. *Journal of Infusion Nursing*, May/June 2005; 28(3).
- Bleasdale, Susan C. et al. Effectiveness of Chlorhexidine Bathing to Reduce Catheter-Associated Bloodstream Infections in Medical Intensive Care Unit Patients. *American Medical Association, Arch Intern Med*, October, 2007; 167(19).
- McWhinney, K. et al. Cardiac Surgical Site Infections at the University of Ottawa Heart Institute: A Case Control Study, Preventative Strategies, and Follow up. *The Canadian Journal of Infection Control*, Winter 1999; 141-147.